



***Dario huli*, a new species of badid from Karnataka, southern India (Teleostei: Percomorpha: Badidae)**

RALF BRITZ¹ & ANVAR ALI²

¹Department of Life Sciences, The Natural History Museum, Cromwell Road, London, SW75BD, United Kingdom.

E-mail: r.britz@nhm.ac.uk

²Conservation Research Group (CRG), Department of Fisheries, St. Albert's College, Kochi, 682 018, India.

E-mail: anvaraliif@gmail.com

Abstract

Dario huli, new species, is described from a small tributary stream of the Tunga River in southern Karnataka, India. It can be distinguished from all its congeners except *D. urops* by the presence of a conspicuous black caudal-fin blotch and by anterior dorsal-fin lappets in males not being produced beyond fin spines. It is readily distinguished from *Dario urops* by the absence of the horizontal suborbital stripe (vs. presence), the presence of a series of up to eight black bars on the body (vs. 2–3 black bars restricted to caudal peduncle), 25 scales in a lateral row (vs. 28), 3–5 tubed lateral-line scales (vs. tubed lateral-line scales completely absent), 13+13=26 vertebrae (vs. 14+14–15=28–29), and the presence of teeth on hypobranchial 3 (vs. absence of teeth).

Key words: taxonomy, freshwater fishes, Western Ghats–Sri Lanka biodiversity hotspot

Introduction

The Western Ghats mountain range along the west coast of Peninsular India harbours a diverse freshwater-fish fauna of close to 300 species (Dahanukar *et al.* 2011). Renewed scientific interest in this fauna in the last few years has led to the discovery of a number of new species whose closest relatives live in the northeast of India and adjacent countries, such as, e.g., the sisorids *Pseudolaguvia austrina* (Radhakrishnan *et al.* 2011) and *P. lapillicola* (Britz *et al.* 2013), the cobitid *Pangio ammophila* (Britz *et al.* 2012a) or the psilorhynchid *Psilorhynchus tenura* (Arunachalam & Muralidharan *et al.* 2008). Another unexpected discovery was the badid *Dario urops* (Britz *et al.* 2012b).

The Badidae comprise a total of 21 valid species (see Kullander & Britz 2002; Geetakumari & Vishwanath 2010; Schindler & Linke 2010; Geetakumari & Kadu 2011; Britz *et al.* 2012) in two genera, *Badis* Bleeker and *Dario* Kullander & Britz. Before the discovery of the southern Indian *D. urops*, the known distribution of the genus *Dario*, with its three included species, was restricted to the Brahmaputra drainage in northern India (*D. dario*) and the Ayeyarwaddy drainage in Myanmar and China (*D. hysginon*, *D. dayingensis*). Recently, an additional species, *D. kajal*, was described from the Meghna river drainage in India (Britz & Kullander 2013).

During fieldwork in southern Karnataka, peninsular India, a species of *Dario* with a distinctive colour pattern was collected from along the banks of a small stream: it is here described as *D. huli*, new species.

Material and methods

Ten measurements and 10 counts were taken following the methods outlined in Kullander & Britz (2002). We adopted Kullander & Britz's (2002) terminology for colour pattern and cephalic lateral-line pores. Counts of vertebrae and procurent rays are based on the single cleared and double-stained (Taylor & Van Dyke, 1985) specimen. Specimens are deposited in the collection of the Bombay Natural History Society (BNHS), Mumbai, India. Information on *D. urops* was taken from the original description (Britz *et al.* 2012b).

Acknowledgements

We are grateful to Nikhil Sood, India Gills, Bangalore, India, who helped to organize the trip during which the new species was discovered. He was a great field companion and instrumental in finding the type locality. We also thank Ravi Rangarajan, Bangalore, India, for his company in the field and for the coordinates of the type locality. Sven Kullander, Swedish Museum of Natural History, Stockholm, Sweden, provided the base map used in figure 3. AA thanks Rajeev Raghavan, Conservation Research Group, Kochi, for financial support through the Mohamed bin Zayed Species Conservation Fund's Project on the 'Lost fishes of the Western Ghats' (MBZSCF 1225670). RB was financially supported by the Life Sciences Department Investment Fund (DIF) of the Natural History Museum, London. Two anonymous reviewers and Rohan Pethiyagoda, Australian Museum, Sydney, provided comments that helped to improve the manuscript. We are also grateful to Rajeev Raghavan, Conservation Research Group, Kochi, for his help during this study.

References

- Arunachalam, M. & Muralidharan, M. (2008) Description of a new species of the genus *Psilorhynchus* (Teleostei: Psilorhynchidae) from a western Ghat stream in southern India. *The Raffles Bulletin of Zoology*, 56, 405–414.
- Britz, R. & Kullander, S.O. (2013) *Dario kajal*, a new species of badid fish from Meghalaya, India (Teleostei: Badidae). *Zootaxa*, 3731 (3), 331–337.
<http://dx.doi.org/10.11646/zootaxa.3731.3.3>
- Britz, R., Ali, A. & Philip, S. (2012) *Dario urops*, a new species of badid fish from the Western Ghats, southern India (Teleostei: Percomorpha: Badidae). *Zootaxa*, 3348, 63–68.
- Britz, R., Ali, A. & Raghavan, R. (2012b). *Pangio ammophila*, a new species of eel-loach from Karnataka, southern India (Teleostei: Cypriniformes: Cobitidae). *Ichthyological Exploration of Freshwaters*, 23, 45–50.
- Britz, R., Ali, A. & Raghavan R. (2013) *Pseudolaguvia lapillicola*, a new species of catfish from Peninsular India (Teleostei: Sisoridae). *Ichthyological Exploration of Freshwaters*, 23, 289–295.
- Dahanukar, N., Raghavan, R., Ali, A., Abraham, R. & Shaji, C.P. (2011) Chapter 3. The status and distribution of freshwater fishes of the Western Ghats. In: S. Molur, S., Smith, K.G., Daniel, B.A. & Darwall, W.R.T. (Compilers) *The status and distribution of freshwater fishes in the Western Ghats, India*. IUCN, Cambridge and Gland, pp. 21–48.
- Geetakumari, K. & Vishwanath, W. (2010) *Badis dibruensis*, a new species (Teleostei: Badidae) from northeastern India. *Journal of Threatened Taxa*, 2, 644–647.
<http://dx.doi.org/10.11609/jott.o2156.644-7>
- Geetakumari, K. & Kadu, K. (2011) *Badis singenensis*, a new fish species (Teleostei: Badidae) from Singen River, Arunachal Pradesh, northeastern India. *Journal of Threatened Taxa*, 3, 2085–2089.
<http://dx.doi.org/10.11609/jott.o2531.2085-9>
- Kullander, S.O. & Britz R. (2002) Revision of the family Badidae (Teleostei: Perciformes), with descriptions of a new genus and ten new species. *Ichthyological Exploration of Freshwaters*, 13, 295–372.
- Pramod, P.K., Fang, F., Rema Devi, K., Liao, T.-Y., Indra, T.J., Jameela Beevi, K.S. & Kullander, S.O. (2010) *Betadevario ramachandrani* a new danionine genus and species from the Western Ghats of India (Teleostei: Cyprinidae: Danioninae). *Zootaxa*, 2519, 31–47.
- Radhakrishnan, K.V., Sureshkumar, S. & Ng, H.H. (2011) *Pseudolaguvia austrina*, a new species of sisorid catfish (Osteichthyes: Siluriformes) from Peninsular India. *Ichthyological Exploration of Freshwaters*, 21, 377–383.
- Taylor, W.R. & Van Dyke, G.G. (1985) Revised procedures for staining and clearing small fishes and other vertebrates for bone and cartilage study. *Cybium*, 9, 107–119.
- Schindler, I. & Linke, H. (2010) *Badis juergenschmidti* - a new species of the Indo-Burmese fish family Badidae (Teleostei: Perciformes) from Myanmar. *Vertebrate Zoology*, 60, 209–216.